

RESEARCH

Open Access

On generating T-X family of distributions using quantile functions

Mohammad A Aljarrah*, Carl Lee and Felix Famoye

* Correspondence:
aljar1ma@cmich.edu
Department of Mathematics, Central
Michigan University, Mt. Pleasant
MI 48859, USA

Abstract

The cumulative distribution function (CDF) of the T - X family is given by $R\{W(F(x))\}$, where R is the CDF of a random variable T , F is the CDF of X and W is an increasing function defined on $[0, 1]$ having the support of T as its range. This family provides a new method of generating univariate distributions. Different choices of the R , F and W functions naturally lead to different families of distributions. This paper proposes the use of quantile functions to define the W function. Some general properties of this T - X system of distributions are studied. It is shown that several existing methods of generating univariate continuous distributions can be derived using this T - X system. Three new distributions of the T - X family are derived, namely, the normal-Weibull based on the quantile of Cauchy distribution, normal-Weibull based on the quantile of logistic distribution, and Weibull-uniform based on the quantile of log-logistic distribution. Two real data sets are applied to illustrate the flexibility of the distributions.

Keywords: Beta-family; Generalized distribution; Survival function; T-X families; Moments

1. Introduction

Statistical distributions are important for parametric inferences and applications to fit real world phenomena. Many methods have been developed to generate statistical distributions in the literature. Some well-known methods in the early days for generating univariate continuous distributions include methods based on differential equations developed by Pearson (1895), methods of translation developed by Johnson (1949), and the methods based on quantile functions developed by Tukey (1960). The interest in developing new methods for generating new or more flexible distributions continues to be active in the recent decades. Lee et al. (2013) indicated that the majority of methods developed after 1980s are the methods of 'combination' for the reason that these new methods are based on the idea of combining two existing distributions or by adding extra parameters to an existing distribution to generate a new family of distributions. A brief summary of some methods in the literature that are related to the method proposed in this article is provided.

McDonald (1984) introduced the generalized beta distributions of the first and second kinds (GB1 and GB2). Subsequently, a further generalization named the generalized beta distribution (GBD) was given by McDonald and Xu (1995), which consists of more than 30 special cases or limiting distributions of GBD, including GB1 and GB2.

Azzalini (1985) introduced a family of skew-normal distributions, SN (λ), defined as $g(x; \lambda) = 2\phi(x)\Phi(\lambda x)$, where ϕ and Φ are probability density function (PDF) and cumulative distribution function (CDF) of $N(0, 1)$, respectively. The skewness is characterized by the parameter λ . For a review of skew-symmetric distributions, one may refer to Kotz and Vicari (2005). Ferreira and Steel (2006) introduced a general framework for generating a family of skewed distributions based on a symmetric distribution. The PDF of the new family has the form

$$g(x|f, p) = f(x)p(F(x)), \tag{1.1}$$

where F is the CDF of a symmetric PDF f and p is a skewed PDF defined on $[0, 1]$.

Marshall and Olkin (1997) proposed a general method for generating a new family of life distributions defined in terms of survival function as

$$\bar{G}(x; \alpha) = \frac{\alpha \bar{F}(x)}{1 - \alpha \bar{F}(x)} = \frac{\alpha \bar{F}(x)}{F(x) + \alpha \bar{F}(x)}, -\infty < x < \infty, \alpha > 0, \tag{1.2}$$

where $\bar{\alpha} = 1 - \alpha$ and $\bar{F} = 1 - F$ is the survival function of the random variable X . For details about life distributions, one may refer to Marshall and Olkin (2010) and Lai (2013).

Eugene et al. (2002) proposed the beta-generated family of distributions, where beta distribution with PDF b is used as the generator. The CDF of the beta generated distribution is defined as $G(x) = \int_0^{F(x)} b(t) dt$, where F is the CDF of any random variable. If X is continuous, the corresponding PDF of the beta generated distribution is

$$g(x) = \frac{f(x)}{B(\alpha, \beta)} F^{\alpha-1}(x)(1-F(x))^{\beta-1}, \alpha > 0, \beta > 0, \tag{1.3}$$

where $B(\alpha, \beta)$ is the beta function. The PDF in (1.3) can be considered as a generalization of the distribution of order statistic (Eugene et al. 2002; Jones 2004). Many researchers have studied the beta generated distributions and their applications by applying different F in Equation (1.3). Examples include Famoye et al. (2004), Akinsete et al. (2008), Cordeiro and Lemonte (2011), and Alshwarbeh et al. (2012).

Jones (2009) and Cordeiro and de Castro (2011) extended the beta-generated family of distributions by using Kumaraswamy distribution $b(t) = \alpha\beta t^{\alpha-1}(1-t)^{\beta-1}$, $t \in (0, 1)$ (Kumaraswamy 1980), instead of the beta distribution. The PDF for Kumaraswamy-generated ($Kw-G$) family of distributions is defined by

$$g(x) = \alpha\beta f(x)F^{\alpha-1}(x)(1-F(x))^{\beta-1}, \alpha > 0, \beta > 0. \tag{1.4}$$

Some examples of the $Kw-G$ distributions are the Kw -Weibull (Cordeiro et al. 2010) and Kw -Gumbel (Cordeiro et al. 2011).

Recently, Alexander et al. (2012) studied the generalized beta- X family by considering $b(t) = cB(a, b)^{-1}t^{ac-1}(1-t)^{b-1}$, $0 < t < 1$, the generalized beta distribution of the first kind introduced by McDonald (1984). The new family is called generalized beta-

generated (GBG) family of distributions. The PDF for GBG family of distributions is given by

$$g(x, \tau, a, b, c) = cB(a, b)^{-1}f(x; \tau)F(x; \tau)^{ac-1}[1-F(x; \tau)]^{b-1}. \quad (1.5)$$

When $c = 1$, the family in (1.5) reduces to the beta- X family in (1.3), and when $a = 1$, (1.5) reduces to the Kw - G family in (1.4).

Alzaatreh et al. (2013b) proposed a general method by replacing the beta PDF with a PDF r of a continuous random variable and applying a function $W(F(x))$ that satisfies some conditions (given in (2.1)) to develop the T - X family. The CDF of the T - X family is defined as

$$G(x) = \int_a^{W(F(x))} r(t) dt = R\{W(F(x))\}, \quad (1.6)$$

where R is the CDF of T . The corresponding PDF (if it exists) of the T - X family of distributions is

$$g(x) = \left\{ \frac{d}{dx} W(F(x)) \right\} r\{W(F(x))\}. \quad (1.7)$$

Different W functions generate different families of T - X distributions. Two continuous distributions of the T - X families that have been studied are Gamma-Pareto distribution (Alzaatreh et al. 2012a) and Weibull-Pareto distribution (Alzaatreh et al. 2013a). When X is discrete, the resulting T - X family is discrete. The T -geometric family generates the discrete analogue to the distribution of any continuous random variable T (Alzaatreh et al. 2012b). For a review of methods for generating univariate continuous distributions, one may refer to Lee et al. (2013).

The T - X family provides a new method to generate distributions by using the function W . A large number of distributions, continuous and discrete, can be generated by applying any two existing univariate distributions based on this method. Alzaatreh et al. (2013b) gave several choices of $W(\lambda)$, including $-\log(1 - \lambda)$, $\lambda/(1 - \lambda)$, $\log(\lambda/(1 - \lambda))$, $\log(-\log \lambda)$. It is clear that there are other choices that can be defined to generate different T - X families. Is there a systematic approach to define the W function for the T - X family? This question will be addressed in this paper.

In Section 2, a method to define the W function for generating T - X families of continuous probability distributions is presented. The W functions defined in Alzaatreh et al. (2013b) are special cases of the general approach. In order to distinguish between the previous T - X family proposed by Alzaatreh et al. (2013b) and the method proposed in this article, we use the abbreviation T - $X(W)$ family for the previous T - X family and the abbreviation T - $X\{Y\}$ for the family defined in Section 2. In Section 3, some properties of the new families are studied. Relationship between the new families and some existing families is given. Also in Section 3, the normal-Weibull distribution based on the quantile function of the Cauchy distribution, the normal-Weibull distribution based on the quantile function of the logistic distribution and Weibull-uniform distribution

based on the quantile function of the log-logistic distribution are defined. Some properties of these three distributions are derived. In Section 4, a general family of life distributions based on survival function using similar methodology of the $T-X\{Y\}$ family is presented. Some properties of the family are investigated. In Section 5, two real data sets are used to illustrate the flexibility of $T-X\{Y\}$ family of distributions. Conclusions are given in Section 6.

2. Generating families of continuous probability distributions using quantile function

The $T-X(W)$ family of distributions in (1.6) is generated by using the function W which satisfies the following conditions (Alzaatreh et al. 2013b):

$$\left. \begin{array}{l} \text{i. } W(F(x)) \in [a, b], \\ \text{ii. } W \text{ is differentiable and monotonically non-decreasing,} \\ \text{iii. } W(F(x)) \rightarrow a \text{ as } x \rightarrow -\infty \text{ and } W(F(x)) \rightarrow b \text{ as } x \rightarrow \infty, \end{array} \right\} \quad (2.1)$$

where $[a, b]$ is the support of the random variable T for $-\infty \leq a < b \leq \infty$.

In this section, a class of W functions wider than the one defined in (2.1) will be considered to define a $T-X$ family. Let $W: (0, 1) \rightarrow (a, b)$, for $-\infty \leq a < b \leq \infty$, be a right-continuous and non-decreasing function such that, $\lim_{\lambda \rightarrow 0^+} W(\lambda) = a$ and $\lim_{\lambda \rightarrow 1^-} W(\lambda) = b$, then the composition $G(x) = R\{W(F(x))\}$, $x \in (-\infty, \infty)$, is a distribution function, because it satisfies the following required conditions for a distribution function:

- (a) G is non-decreasing,
- (b) G is right-continuous,
- (c) $G(x) \rightarrow 0$ as $x \rightarrow -\infty$ and $G(x) \rightarrow 1$ as $x \rightarrow \infty$.

If T has a PDF r with support (a, b) , then

$$G(x) = \int_a^{W(F(x))} r(t) dt. \quad (2.2)$$

Note that if both functions W and F are absolutely continuous, then G in (2.2) is absolutely continuous and has a density function $g(x) = \frac{d}{dx}G(x)$.

A general method to define W function for generating $T-X$ families is now proposed. It is assumed that the random variable T has support on the interval (a, b) . Let P be the CDF of the random variable Y taking values on (a, b) , and define the quantile function of the distribution P by

$$Q_Y(\lambda) = \inf\{y : P(y) \geq \lambda\}, \quad \lambda \in (0, 1).$$

If P is continuous and strictly increasing then $Q_Y = P^{-1}$ is continuous and strictly increasing (Shorack and Wellner 1986). We take W to be the quantile function of a strictly increasing distribution function P for the random variable Y , namely, $W(\lambda) = Q_Y(\lambda)$, $\lambda \in (0, 1)$, then Q_Y is continuous and non-decreasing, and the CDF of a $T-X\{Y\}$ family using the quantile function Q_Y is defined as

$$G(x) = \int_a^{Q_Y(F(x))} r(t) dt = R\{Q_Y(F(x))\}, \quad x \in (-\infty, \infty). \quad (2.3)$$

If we assume further that Y has a density $p(y) > 0$ for all y in a neighborhood of $Q_Y(\lambda)$ where $\lambda \in (0, 1)$, then $\frac{d}{d\lambda}Q_Y(\lambda)$ exists and equals $[p(Q_Y(\lambda))]^{-1}$ (Shorack and Wellner 1986), and hence the corresponding PDF associated with (2.3) is

$$g(x) = \frac{f(x)}{p\{Q_Y(F(x))\}} r\{Q_Y(F(x))\}. \tag{2.4}$$

Note that the PDF defined in (2.4) can be easily used to generate a $T-X\{Y\}$ family of distributions by applying the quantile function of any existing distribution.

The notation X sometimes represents the random variable with PDF f and sometimes represents the random variable with PDF g . Where there may be confusion, the notations X_f for the random variable X with PDF f and X_g for the random variable X with PDF g are used. The term moment refers to non-central moment, unless otherwise specified.

Lemma 1:

- (a) If the random variables X_f and Y have the same distribution with the same parameters, then $G = R$.
- (b) If the random variables T and Y have the same distribution with the same parameters, then $G = F$.

Proof: The proofs of (a) and (b) follow from definition (2.4). □

Some properties of the $T-X\{Y\}$ family:

1. Any PDF f can be represented as the PDF defined in (2.4) by considering $Q_Y = R^{-1}$.
2. The support of the new random variable defined in (2.4) is the same as the support of the random variable with PDF f .
3. If the support of the random variable Y is $[c, d]$ with $[a, b] \subset [c, d]$, then the PDF in (2.4) is defined with support $[F^{-1}(P(a)), F^{-1}(P(b))]$.
4. The relationship between the random variable X_g with PDF in (2.4) and T is given by $T = Q_Y(F(X_g))$ and hence, $X_g = F^{-1}(P(T))$ when F^{-1} exists, where P is the CDF of Y with the corresponding quantile function Q_Y . Using this relation, one can generate the random variable X_g by generating the random variable T and then computing $X_g = F^{-1}(P(T))$. Similarly, one can compute the moments of X_g by using $E(X_g^n) = E\{[F^{-1}(P(T))]^n\}$.
5. The hazard function, $h_g(x) = g(x)/\bar{G}(x)$, for the random variable X_g in (2.4) is given by $h_g(x) = \frac{f(x)}{p\{Q_Y(F(x))\}} h_r\{Q_Y(F(x))\}$, where h_r is the hazard function for the random variable T with PDF r .
6. The quantile function of the new random variable in (2.4) is given by $Q_{X_g}(\lambda) = Q_{X_f}\{P(Q_T(\lambda))\}$, $\lambda \in (0, 1)$, where Q_{X_f} and Q_T are the quantile functions of the random variables X_f and T respectively.

3. Some $T-X\{Y\}$ families and properties

3.1 Some $T-X\{Y\}$ families based on different quantile functions

The quantile function of a random variable Y may not be explicitly represented. However, many of the existing continuous random variables are one-to-one functions and they have explicit quantile functions. The quantile functions of these random variables

can be used to generate new $T-X\{Y\}$ families. The following example illustrates how to derive the $T-X\{Y\}$ family of distributions.

Example: $T-X\{\log\text{-logistic}\}$ family:

Let the random variable Y follow the log-logistic distribution with parameters α and β . The PDF and quantile function are, respectively, $p(y) = (\beta/\alpha)(y/\alpha)^{\beta-1}/(1+(y/\alpha)^\beta)^2$, $y \geq 0$, and $Q_Y(\lambda) = \alpha(\lambda/(1-\lambda))^{1/\beta}$, $\lambda \in (0, 1)$. Therefore, $p(Q_Y(\lambda)) = (\beta/\alpha)\lambda^{(\beta-1)/\beta}(1-\lambda)^{(\beta+1)/\beta}$, and the definition in (2.4) gives the PDF of $T-X\{\log\text{-logistic}\}$ family as

$$g(x) = \frac{(\alpha/\beta)f(x)}{F^{(\beta-1)/\beta}(x)(1-F(x))^{(\beta+1)/\beta}} r\left\{\alpha\left(\frac{F(x)}{1-F(x)}\right)^{1/\beta}\right\}. \tag{3.1}$$

When $\alpha = \beta = 1$, the family in (3.1) reduces to $g(x) = \frac{f(x)}{(1-F(x))^2} r\{F(x)/(1-F(x))\}$. This PDF can be written in terms of hazard and survival functions of X_f as $g(x) = \frac{h_f(x)}{\bar{F}(x)} r\{(1-\bar{F}(x))/\bar{F}(x)\}$, where h_f is the hazard function and \bar{F} is the survival function of the random variable X_f .

Table 1 lists the probability density functions of some $T-X\{Y\}$ families based on different quantile functions. Each family g is based on a given quantile function that defines many subfamilies of distributions.

Table 1 $T-X\{Y\}$ families based on different quantile functions

Random variable Y and support of $r(t)$	The quantile function $Q_Y(\lambda)$	Family of probability density function $g(x)$ defined in (2.4)
Exponential $(0, \infty)$	$-b \log(1-\lambda), b > 0$	$\frac{bf(x)}{1-F(x)} r\{-b \log(1-F(x))\}$
Weibull $(0, \infty)$	$\gamma[-\log(1-\lambda)]^{1/c}, \gamma, c > 0$	$\frac{\gamma f(x) r\{\gamma[-\log(1-F(x))]^{1/c}\}}{c(1-F(x))\{-\log(1-F(x))\}^{(c-1)/c}}$
Rayleigh $(0, \infty)$	$\{-2b^2 \log(1-\lambda)\}^{1/2}, b > 0$	$\frac{bf(x) r\{\{-2b^2 \log(1-F(x))\}^{1/2}\}}{(1-F(x))\{-2 \log(1-F(x))\}^{1/2}}$
Dagum $(0, \infty)$	$\beta\left\{\frac{\lambda^{1/p}}{1-\lambda^{1/p}}\right\}^{1/a}, a, \beta, p > 0$	$\frac{\beta f(x) r\{\beta F^{1/ap}(x)/(1-F^{1/p}(x))^{1/a}\}}{apF^{1-1/ap}(x)(1-F^{1/p}(x))^{1+1/a}}$
Lomax $(0, \infty)$	$\frac{1}{a}\left\{\frac{1-(1-\lambda)^{1/k}}{(1-\lambda)^{1/k}}\right\}, a, k > 0$	$\frac{f(x)}{ka(1-F(x))^{1/k+1}} r\left\{\frac{1-(1-F(x))^{1/k}}{a(1-F(x))^{1/k}}\right\}$
Log-logistic $(0, \infty)$	$\alpha\left(\frac{\lambda}{1-\lambda}\right)^{1/\beta}, \alpha, \beta > 0$	$\frac{af(x) r\{\alpha(F(x)/(1-F(x)))^{1/\beta}\}}{\beta F^{(\beta-1)/\beta}(x)(1-F(x))^{(\beta+1)/\beta}}$
Exponentiated Exponential $(0, \infty)$	$-\frac{1}{\theta} \log(1-\lambda^{1/a}), \theta, a > 0$	$\frac{f(x) r\{-(1/\theta) \log(1-F^{1/a}(x))\}}{a\theta F^{(a-1)/a}(x)(1-F^{1/a}(x))}$
Cauchy $(-\infty, \infty)$	$a + b \{\tan(\pi(\lambda - 0.5))\}, b > 0$	$\frac{\pi b f(x) r\{a+b \tan\{\pi(F(x)-1/2)\}\}}{\cos^2\{\pi(F(x)-1/2)\}}$
Extreme value (Gumbel) $(-\infty, \infty)$	$a - b \log(-\log \lambda), b > 0$	$\frac{bf(x) r\{a-b \log(-\log F(x))\}}{-F(x) \log F(x)}$
Laplace $(-\infty, \infty)$	$\begin{cases} a + b \log(2\lambda), & \lambda < 0.5 \\ a - b \log(2(1-\lambda)), & \lambda \geq 0.5 \end{cases}$ $a, b > 0$	$\begin{cases} \frac{bf(x) r\{a + b \log(2F(x))\}}{F(x)}, & F(x) < .5 \\ \frac{bf(x) r\{a - b \log\{2(1-F(x))\}}}{1-F(x)}, & F(x) \geq .5 \end{cases}$
Logistic $(-\infty, \infty)$	$a + b \log\left(\frac{\lambda}{1-\lambda}\right), b > 0$	$\frac{bf(x) r\{a+b \log(F(x)/(1-F(x)))\}}{F(x)(1-F(x))}$
Generalized logistic (II) $(-\infty, \infty)$	$\log\left\{\frac{1-(1-\lambda)^{1/a}}{(1-\lambda)^{1/a}}\right\}, a > 0$	$\frac{f(x) r\{\log\{1-(1-F(x))^{1/a}\}/(1-F(x))^{1/a}\}}{a(1-F(x))\{1-(1-F(x))^{1/a}\}}$

Common supports of random variables X_f and T are $[0, 1]$, $(0, \infty)$, or $(-\infty, \infty)$. Beta- X , $Kw-G$ and GBG are $T-X\{uniform\}$ families with T being defined on $[0, 1]$. The W functions given in Alzaatreh et al. (2013b) can be defined by the quantile functions of random variable Y as follows: $W(\lambda) = -\log(1 - \lambda)$, $\lambda \in (0, 1)$, is the quantile function of standard exponential distribution, $W(\lambda) = \lambda/(1 - \lambda)$ is the quantile function of log-logistic distribution with parameters $\alpha = \beta = 1$, and $W(\lambda) = \log(\lambda/(1 - \lambda))$ is the quantile function of logistic distribution with scale parameter $b = 1$ and location parameter $a = 0$. Many other W functions can be defined by using the quantile function approach. The $T-X(W)$ families defined in Alzaatreh et al. (2013b) derive their parameters from the random variables T and X and none from the W function. The $T-X(W)$ can be derived through the $T-X\{Y\}$ framework by noting that the W function is the quantile function for the random variable Y . One advantage of using the $T-X\{Y\}$ framework is that one can keep one or more parameters from the distribution of Y . In particular, keeping a shape parameter from Y can add more flexibility to the new distribution.

3.2 Some properties of $T-X\{Y\}$ families

In the following, we assume (if necessary) the mentioned expectations exist and are finite.

Theorem 1: Let X_f be a non-negative random variable with PDF $f(x)$, and let $E(X_f^n)$ denote the n^{th} moment of X_f , then

$$E(X_g^n) \leq E(X_f^n) \cdot E(\{\bar{P}(T)\}^{-1}),$$

where $E(X_g^n)$ is the n^{th} moment of the random variable with density in (2.4), $\bar{P} = 1 - P$ is the survival function of the CDF P , and T is the random variable with PDF r .

Proof: By definition,

$$\begin{aligned} E(X_f^n) &= \int_0^\infty x^n f(x) dx \geq \int_{F^{-1}\{P(t)\}}^\infty x^n f(x) dx \geq (F^{-1}\{P(t)\})^n \int_{F^{-1}\{P(t)\}}^\infty f(x) dx \\ &= (F^{-1}\{P(t)\})^n \cdot \bar{P}(t). \end{aligned}$$

Hence,

$$(F^{-1}\{P(t)\})^n \leq E(X_f^n) \{\bar{P}(t)\}^{-1}. \tag{3.2}$$

Using property (4) in Section 2 and (3.2) yields

$$E(X_g^n) = E\left([F^{-1}\{P(T)\}]^n\right) \leq E(X_f^n) \cdot E(\{\bar{P}(T)\}^{-1}). \quad \square$$

The entropy of a random variable is a measure of the variation of uncertainty. Shannon's (1948) entropy of the random variable X with density g is defined as $E\{-\log(g(X))\}$.

Theorem 2: Shannon's entropy η_{X_g} of X_g is given by

$$\eta_{X_g} = E\left\{\log q_{X_f}(P(T))\right\} + E\{\log p(T)\} + \eta_T,$$

where η_T is Shannon's entropy for the random variable T with PDF r , and q_{X_f} is the quantile density function of X_f .

Proof: By definition,

$$\begin{aligned} \eta_{X_g} &= E\{-\log(g(X_g))\} \\ &= -E\{\log f(X_g)\} + E\{\log p(Q_Y\{F(X_g)\})\} - E\{\log r(Q_Y\{F(X_g)\})\}. \end{aligned}$$

Note that the random variable $T = Q_Y\{F(X_g)\}$ has the PDF r , and $X_g = F^{-1}\{P(T)\}$. Thus,

$$\begin{aligned} \eta_{X_g} &= -E\{\log f(F^{-1}\{P(T)\})\} + E\{\log p(T)\} + \eta_T \\ &= E\{\log q_{X_f}(P(T))\} + E\{\log p(T)\} + \eta_T. \quad \square \end{aligned}$$

3.3 Relationship between $T-X\{Y\}$ family and some existing families of distributions

Many existing families of distributions can be generated by using the quantile function approach defined in (2.4). Four examples are given in the following.

Generalized beta-generated (GBG) family introduced by Alexander et al. (2012):

The GBG family is the generalized beta- $X\{\text{uniform}\}$ family. It can also be derived as follows: By setting $\alpha = \beta = 1$ and $p = 1/c$ in the Dagum quantile function in Table 1, the PDF of the $T-X\{\text{Dagum}\}$ family is given by

$$g(x) = \frac{cf(x)r\{F^c(x)/(1-F^c(x))\}}{F^{1-c}(x)(1-F^c(x))^2}. \quad (3.3)$$

By taking r in (3.3) to be $r(t) = \{B(a, b)\}^{-1}t^{a-1}(1+t)^{-b-a}$, $t > 0$, which is the PDF of the inverted beta random variable, the family of probability density functions is obtained as

$$g(x) = cB(a, b)^{-1}f(x)F^{ca-1}(x)(1-F^c(x))^{b-1}, a, b > 0. \quad (3.4)$$

The family in (3.4) is the generalized beta-generated (GBG) family in (1.5).

Family of skewed distributions defined in Ferreira and Steel (2006):

The family of skewed distributions in (1.1) defined by Ferreira and Steel (2006) can be represented in the form of $T-X\{Y\}$ system by considering the quantile function of a standard uniform distribution, $Q_Y(\lambda) = \lambda$, where F is the CDF of a symmetric PDF f and the r is a skewed PDF having support $[0, 1]$.

$T-X$ family defined in Alzaatreh et al. (2013b):

Alzaatreh et al. (2013b) studied the $T-X(W)$ family using $W(F(x)) = -\log(1 - F(x))$. By using the quantile function approach, let $\lambda = F(x)$, then $W(\lambda) = -\log(1 - \lambda)$ is the quantile function of standard exponential distribution. Hence, the $T-X$ family studied by Alzaatreh et al. (2013b) is the $T-X\{\text{exponential}\}$. According to the authors, the $T-X\{\text{exponential}\}$ is a family of distributions arising from the hazard function of X_f . If h_f is the hazard function and H_f is the cumulative hazard function of the random variable X_f , and the exponential distribution has mean b , the PDF of the $T-X\{\text{exponential}\}$ family is $g(x) = bh_f(x)r\{bH_f(x)\}$. In a similar way, $T-X\{\text{Weibull}\}$ and $T-X\{\text{Rayleigh}\}$ families can be considered as families of distributions arising from hazard functions of X_f . The PDF of the $T-X\{\text{Weibull}\}$ family is $g(x) = (\gamma/c)h_f(x)r\{\gamma(H_f(x))^{1/c}\}/(H_f(x))^{(c-1)/c}$ and the PDF of $T-X\{\text{Rayleigh}\}$ is $g(x) = bh_f(x)r\{(2b^2H_f(x))^{1/2}\}/(2H_f(x))^{1/2}$.

Generalized beta distribution introduced by McDonald and Xu (1995):

Setting $\alpha = \beta = 1$ in the PDF of the $T-X\{\text{log-logistic}\}$ family in (3.1) and taking $r(t) = \{B(p, q)\}^{-1} t^{p-1} (1+t)^{-p-q}$, $t > 0$, the PDF of the inverted beta random variable with parameters p and q , yields

$$g(x) = \frac{f(x)}{B(p, q)} F^{p-1}(x) (1-F(x))^{q-1}, p, q \geq 0. \tag{3.5}$$

Note that (3.5) is the beta-generated family. By taking $F(x) = (x/b)^a / (1 + c(x/b)^a)$, which is the CDF of a truncated log-logistic distribution with $0 < x^a < b^a / (1 - c)$, $0 \leq c < 1$, a and b positive in (3.5), the generalized beta distribution introduced by McDonald and Xu (1995) is obtained as

$$GB(x; a, b, c, p, q) = \frac{ax^{ap-1} (1 - (1-c)(x/b)^a)^{q-1}}{b^{ap} B(p, q) (1 + c(x/b)^a)^{p+q}}, \text{ for } 0 < x^a < \frac{b^a}{1-c},$$

with $0 \leq c < 1$, and a, b, p and q positive.

Taking $F(x)$ in (3.5) to be $F(x) = e^{(x-\delta)/\sigma} / (1 + ce^{(x-\delta)/\sigma})$, which is the CDF of a truncated logistic distribution with $-\infty < (x - \delta)/\sigma < \ln(1/(1 - c))$, $0 \leq c < 1$ and $\sigma > 0$, yields

$$g(x) = e^{p(x-\delta)/\sigma} (1 - (1-c)e^{(x-\delta)/\sigma})^{q-1} / \left[\sigma B(p, q) (1 + ce^{(x-\delta)/\sigma})^{p+q} \right]. \tag{3.6}$$

The PDF in (3.6) is the exponential generalized beta distribution in McDonald and Xu (1995).

3.4 Three examples of new distributions derived from $T-X\{Y\}$ family

Table 1 contains many $T-X$ families based on different quantile functions. Three new distributions, normal-Weibull{Cauchy}, normal-Weibull{logistic} and Weibull-uniform {log-logistic} distributions are introduced, and some properties of these distributions are studied.

Normal-Weibull{Cauchy} distribution:

Setting $a = 0$ and $b = 1$ in the $T-X\{\text{Cauchy}\}$ family in Table 1, and letting r be $N(\mu, \sigma^2)$, the normal- $X\{\text{Cauchy}\}$ sub-family is given by


$$g(x) = \frac{\sqrt{\pi} f(x)}{\sigma \sqrt{2} \cos^2\{\pi(F(x)-1/2)\}} \exp\left\{ \frac{-1}{2} \left(\frac{\tan\{\pi(F(x)-1/2)\} - \mu}{\sigma} \right)^2 \right\}. \tag{3.7}$$

Substituting $F(x) = 1 - \exp\{-(x/\gamma)^c\}$, the CDF of Weibull distribution in (3.7) yields

$$g(x) = \frac{\sqrt{\pi}(c/\gamma)(x/\gamma)^{c-1} \exp\{-(x/\gamma)^c\}}{\sigma \sqrt{2} \cos^2\{\pi(1/2 - \exp\{-(x/\gamma)^c\})\}} \exp\left\{ - \left(\frac{\tan\{\pi(1/2 - \exp\{-(x/\gamma)^c\})\} - \mu}{\sigma \sqrt{2}} \right)^2 \right\}, \tag{3.8}$$

for $x > 0$, and $\sigma, c, \gamma > 0$. The random variable with the PDF in (3.8) is said to follow a four-parameter normal-Weibull{Cauchy} (NW{C}) distribution. A location parameter δ can be included in (3.8) by writing x as $(x - \delta)$ leading to a five-parameter distribution.

Plots of the NW{C} density function for different parameter values are given in Figure 1. The graphs in Figure 1 show that the NW{C} distribution can be right skewed, left skewed, unimodal or bimodal.



Lemma 2: The n^{th} moment of the NW{C} random variable with PDF in (3.8) exists for any $\mu, \sigma > 0, c > 0, \gamma > 0$ and satisfies the inequality

$$E\left(X_g^n\right) \leq \gamma^n \Gamma(1+n/c) \left\{ 4\Phi(1) + \frac{3\sqrt{\pi}\sigma}{2\sqrt{2}} \exp\left(-\frac{(1-\mu)^2}{2\sigma^2}\right) + \left(\frac{3\pi\mu}{2}\right)(1-\Phi(1)) \right\}, \quad (3.9)$$

where Φ is the CDF of a normal distribution with parameters μ and σ .

Proof: The n^{th} moment for Weibull random variable is $E\left(X_f^n\right) = \gamma^n \Gamma(1+n/c)$. The CDF of standard Cauchy distribution is $P(y) = 1/2 + (1/\pi)\tan^{-1}(y)$, so $1 - P(T) = 1/2 - (1/\pi)\tan^{-1}(T)$, where $-\infty < T < \infty$. When $T \leq 1$, $1 - P(T) \geq 1/4$ and hence $(1 - P(T))^{-1} \leq 4$. When $T > 1$ and by using the series $\tan^{-1}(T) = \frac{\pi}{2} + \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n-1)T^{2n-1}}$ (Polyanin and Manzhirov 2008),

$$\begin{aligned} 1/2 - (1/\pi) \tan^{-1}(T) &= \frac{1}{2} - \frac{1}{\pi} \tan^{-1}(T) \\ &= \frac{1}{\pi} \left\{ \frac{1}{T} - \frac{1}{3T^3} + \frac{1}{5T^5} - \frac{1}{7T^7} + \dots + \frac{(-1)^{n+1}}{(2n-1)T^{2n-1}} \dots \right\} \\ &> \frac{1}{\pi} \left\{ \frac{1}{T} - \frac{1}{3T^3} \right\} = \frac{1}{\pi} \left\{ \frac{3T^2 - 1}{3T^3} \right\} = \frac{1}{\pi} \left\{ \frac{2T^2 + T^2 - 1}{3T^3} \right\} > \frac{2}{3\pi T}. \end{aligned}$$

Hence, $(1 - P(T))^{-1} < (3\pi/2)T$. Since the random variable T with PDF r has a normal distribution with parameters μ and σ , then by using Theorem 1

$$\begin{aligned} E\left(X_g^n\right) &\leq E\left(X_f^n\right) E\left(\{1 - P(T)\}^{-1}\right) \leq \gamma^n \Gamma(1+n/c) \left(\int_{-\infty}^1 4r(t)dt + (3\pi/2) \int_1^{\infty} t \cdot r(t)dt \right), \\ &= \gamma^n \Gamma(1+n/c) \left(\int_{-\infty}^1 4r(t)dt + (3\pi\sigma/2) \int_1^{\infty} \frac{t-\mu}{\sigma} \cdot r(t)dt + (3\pi/2) \int_1^{\infty} \mu \cdot r(t)dt \right), \end{aligned}$$

and the result in (3.9) follows. \square

Normal-Weibull{logistic} distribution:

Setting $a = 0$ and $b = 1$ in the PDF of the $T-X\{logistic\}$ family in Table 1, and taking r to be $N(\mu, \sigma)$, the PDF of normal distribution, and F to be the CDF of Weibull distribution, $F(x) = 1 - \exp\{-(x/\gamma)^c\}$, the normal-Weibull{logistic} (NW{L}) distribution is obtained as

$$g(x) = \frac{c(x/\gamma)^{c-1} \exp\{(x/\gamma)^c\}}{\gamma\sigma\sqrt{2\pi}(\exp\{(x/\gamma)^c\}-1)} \exp\left\{-\frac{1}{2}\left(\frac{\log\{\exp(x/\gamma)^c-1\}-\mu}{\sigma}\right)^2\right\}, x > 0, \sigma, c, \gamma > 0.$$

Plots of NW{L} density function for different parameter values are given in Figure 2. The graphs in Figure 2 show that the NW{L} distribution can be reversed J-shape, skewed to the right or skewed to the left or bimodal.

Lemma 3: The n^{th} moment of the NW{L} random variable exists for any $\sigma > 0, c > 0, \gamma > 0$ and satisfies the inequality

$$E\left(X_g^n\right) \leq \gamma^n \Gamma(1 + n/c) (1 + \exp\{\mu + 0.5\sigma^2\}). \tag{3.10}$$

Proof: The n^{th} moment for the Weibull random variable is $E(X_f^n) = \gamma^n \Gamma(1 + n/c)$. The CDF of standard logistic random variable is $P(y) = \exp(y)/(1 + \exp(y))$. Since the random variable T has normal distribution with parameters μ and σ , then $E(\{1 - P(T)\}^{-1}) = E(1 + \exp(T)) = 1 + \exp(\mu + 0.5\sigma^2)$. By using Theorem 1, the result in (3.10) follows. \square

Weibull-uniform{log-logistic} distribution:

Setting $\alpha = \beta = 1$ in the PDF of the $T-X\{log-logistic\}$ family in (3.1), and taking r to be the PDF of Weibull distribution, $r(t) = (c/\gamma)(x/\gamma)^{c-1} \exp\{-(x/\gamma)^c\}$ and F to be the CDF of uniform distribution, $F(x) = (x - a)/(b - a)$, the Weibull-uniform{log-logistic} (WU{LL}) distribution is obtained as

$$g(x) = \frac{c(b-a)}{\gamma(b-x)^2} \left(\frac{x-a}{\gamma(b-x)}\right)^{c-1} \exp\left\{-\left(\frac{x-a}{\gamma(b-x)}\right)^c\right\}, a < x < b, c, \gamma > 0.$$




Figure 2 The PDF of Normal-Weibull{logistic} for various values of μ, σ, c and γ .

Plots of $WU\{LL\}$ density function for different parameter values are given in Figure 3. The graphs in Figure 3 show that the $WU\{LL\}$ distribution can be reversed J-shape, skewed to the right or skewed to the left or bimodal.

Lemma 4: The n^{th} moment of the $WU\{LL\}$ random variable exists for any $b > a, c > 0, \gamma > 0$ and satisfies the inequality

$$E(X_g^n) \leq \frac{b^{n+1} - a^{n+1}}{(n+1)(b-a)} (1 + \gamma \Gamma(1 + 1/c)). \tag{3.11}$$

Proof: The n^{th} moment for the uniform random variable is $E(X_f^n) = \frac{b^{n+1} - a^{n+1}}{(n+1)(b-a)}$. The CDF of standard log-logistic random variable is $P(y) = y/(1+y)$. Since the random variable T has Weibull distribution with parameters c and γ , then $E\{(1 - P(T))^{-1}\} = E(1 + T) = 1 + \gamma \Gamma(1 + 1/c)$. By using Theorem 1, the result in (3.11) follows. \square

4. The family of $T-X\{Y\}$ distributions based on survival functions

Instead of using the CDF F in (2.2), one can use the survival function \bar{F} and apply similar method to generate a new family of distributions in terms of survival functions.

If \bar{P} and Q_Y are the survival and quantile functions of the random variable Y , then $\bar{P}^{-1}(F(x)) = Q_Y(1 - F(x)) = Q_Y(\bar{F}(x))$. A new family $T-X\{Y\}$ of distributions in terms of the survival function of X is defined as

$$\bar{G}(x) = \int_a^{\bar{P}^{-1}(F(x))} r(t) dt = \int_a^{Q_Y(\bar{F}(x))} r(t) dt = R\{Q_Y(\bar{F}(x))\}. \tag{4.1}$$

The corresponding PDF associated with (4.1) is

$$g(x) = \frac{f(x)}{p\{Q_Y(\bar{F}(x))\}} r\{Q_Y(\bar{F}(x))\}. \tag{4.2}$$

The family of life distributions introduced by Marshall and Olkin (1997) can be derived using (4.1) as follows. By using $Q_Y(\lambda) = a(\lambda/(1-\lambda))^{1/b}$, the quantile function




Figure 3 The PDF of Weibull-uniform{log-logistic} for various values of c and γ when $a = 0$ and $b = 5$.

of the log-logistic distribution and $R(t) = \eta t / (1 + \eta t)$, the CDF of log-logistic distribution with scale parameter $1/\eta$ in (4.1) can be written as

$$\bar{G}(x) = \frac{\eta \left\{ a [\bar{F}(x) / (1 - \bar{F}(x))]^{1/b} \right\}}{1 + \eta \left\{ a [\bar{F}(x) / (1 - \bar{F}(x))]^{1/b} \right\}} = \frac{\eta a \bar{F}^{1/b}(x)}{F^{1/b}(x) + \eta a \bar{F}^{1/b}(x)}. \tag{4.3}$$

Letting $\eta a = \alpha$ and $1/b = \beta$, (4.3) becomes $\bar{G}(x) = \frac{\alpha \bar{F}^\beta(x)}{F^\beta(x) + \alpha \bar{F}^\beta(x)}$, which reduces to Marshall-Olkin's family in (1.2) when $\beta = 1$.

The following theorem gives the relation between the moments of the random variables defined in (2.4) and (4.2) when the PDF f is symmetric.

Theorem 3: Let $E_1(X_g^n)$ and $E_2(X_g^n)$ denote the n^{th} moments of the random variables in (2.4) and (4.2) respectively. If f is symmetric, then

$$E_2(X_g^n) = \sum_{i=0}^n (-1)^i \binom{n}{i} (2m_f)^{n-i} E_1(X_g^i),$$

where m_f is the median of the random variable X_f with PDF f .

Proof: The n^{th} moment of (2.4) and (4.2) are $E_1(X_g^n) = \int_0^1 \frac{\{F^{-1}(v)\}^n}{p\{Q_Y(v)\}} r\{Q_Y(v)\} dv$ and $E_2(X_g^n) = \int_0^1 \frac{\{F^{-1}(1-u)\}^n}{p\{Q_Y(u)\}} r\{Q_Y(u)\} du$, where the substitutions $v = F(x)$ and $u = 1 - F(x)$ are applied to (2.4) and (4.2) respectively. When f is symmetric, $F^{-1}(1 - u) = 2m_f - F^{-1}(u)$, $u \in [0, 1]$. By using the binomial theorem,

$$\begin{aligned} E_2(X_g^n) &= \int_0^1 \frac{\{F^{-1}(1-u)\}^n}{p\{Q_Y(u)\}} r\{Q_Y(u)\} du \\ &= \sum_{i=0}^n (-1)^i \binom{n}{i} (2m_f)^{n-i} \int_0^1 \frac{\{F^{-1}(u)\}^i}{p\{Q_Y(u)\}} r\{Q_Y(u)\} du, \\ &= \sum_{i=0}^n (-1)^i \binom{n}{i} (2m_f)^{n-i} E_1(X_g^i). \end{aligned} \quad \square$$

Theorem 4: Let X_f be non-negative random variable with PDF f , and let $E(X_f^n)$ denote the n^{th} moment of X_f , then

$$E(X_g^n) \leq E(X_f^n) \cdot E(\{P(T)\}^{-1}),$$

where $E(X_g^n)$ is the n^{th} moment of the random variable with density in (4.2), and T is the random variable with PDF r .

Proof: The proof is similar to the proof of Theorem 1 after noting that the relation between the random variables X_g in (4.2) and T is given by $X_g = F^{-1}(\bar{P}(T))$. \square

Similar to Theorem 2, Shannon's entropy η_{X_g} for random variable X_g with PDF in (4.2) is given in the following theorem.

Theorem 5: Let X_g be a random variable with density in (4.2). Shannon's entropy η_{X_g} is given by $\eta_{X_g} = E\left\{ \log q_{X_f}(\bar{P}(T)) \right\} + E\{ \log p(T) \} + \eta_T$.

Proof: The proof is similar to that of Theorem 2. \square

5. Application

In this section, we apply the $NW\{C\}$ distribution to fit two data sets. The first data is the famous Old Faithful Geyser eruption data ($n = 272$) obtained from Härdle (1991, p. 201). The data is the duration time of eruption (in minutes) taken during August 1st to August 15th, 1985 (Dekking et al. 2005). The second data set is USS Halfbeak diesel engine data ($n = 71$) studied by Ascher and Feingold (1984, p. 75) and Meeker and Escobar (1998, p. 415). The data is the time of unscheduled maintenance actions for the USS Halfbeak number 4 main propulsion diesel engine over 25.518 operating hours (Meeker and Escobar 1998, p. 415).

5.1 The famous old faithful Geyser eruption data

As shown in Figure 4, the data has two distinct modes. A common approach for fitting such a bimodal data is by using mixture distributions. Arellano-Valle et al. (2010) applied flexible epsilon-skew-normal distribution to fit the data and their fit is the same as that of mixture-normal distribution. Four distributions, a four-parameter $NW\{C\}$ in (3.8), a five-parameter $NW\{C\}$, mixture normal, and beta-normal are applied to fit the data using maximum likelihood technique. Table 2 contains the estimates, standard errors of the estimates, log-likelihood values, AIC, K-S test statistics and the corresponding p -values.

The results in Table 2 indicate that the five-parameter $NW\{C\}$ provides the best fit followed by mixture normal based on all three measures, log-likelihood, AIC and K-S statistic. When bimodality is a population characteristic, it may be more appropriate to fit the data with one distribution, instead of fitting the data by the mixture of two distributions. The $NW\{C\}$ distribution can fit well a wide variety of distribution shapes, including bimodal data such as Old Faithful Geyser eruption data. Figure 4 displays the estimated PDF of the distributions that provide adequate fit to the data.

From Figure 4, the addition of the fifth parameter, which is a measure of location, has some effect on the fit. By using either a likelihood ratio test or the Wald test for the significance of the parameter δ , we observe that the parameter is significantly different from zero. Thus, a five-parameter $NW\{C\}$ (and not a four-parameter $NW\{C\}$) should be used to fit the bivariate data. According to Johnson et al. (1994, p. 12), four-parameter




Table 2 The MLEs and goodness-of-fit statistics for the famous Old Faithful Eruption data

	Five-parameter NW{C}	Mixture-normal ^a	Four-parameter NW{C}	Beta-normal
MLE (SE ^b)	$\hat{\mu} = 1.883 (0.447)$ $\hat{\sigma} = 5.156 (0.645)$ $\hat{c} = 2.032 (0.150)$ $\hat{\gamma} = 1.885 (0.116)$ $\hat{\delta} = 1.343 (0.060)$	$\hat{\mu}_1 = 4.273 (0.034)$ $\hat{\sigma}_1 = 0.437 (0.027)$ $\hat{\mu}_2 = 2.019 (0.026)$ $\hat{\sigma}_2 = 0.236 (0.023)$ $\hat{\rho} = 0.652 (0.029)$	$\hat{\mu} = 0.644 (0.288)$ $\hat{\sigma} = 3.756 (0.438)$ $\hat{c} = 3.899 (0.168)$ $\hat{\gamma} = 3.623 (0.031)$	$\hat{\mu} = 3.163 (0.007)$ $\hat{\sigma} = 0.238 (0.001)$ $\hat{a} = 0.083 (0.007)$ $\hat{b} = 0.060 (0.004)$
Log-likelihood	-270.0	-276.35	-292.3	-372.566
AIC	550.0	562.7	592.6	753.1
K-S statistic	0.042	0.049	0.075	0.151
p-value	0.712	0.539	0.096	8.908e-06

^aMixture normal is defined as $pN(\mu_1, \sigma_1) + (1 - p)N(\mu_2, \sigma_2)$; ^bSE: standard error of the MLE.

distributions should be sufficient for most practical purposes. The authors went on to state and we quote "... but it is doubtful whether the improvement obtained by including a fifth or sixth parameter is commensurate with the extra labor involved". For this application, adding the fifth parameter to the NW{C} improves the fit with an increase of more than 22 points in the log-likelihood value. Furthermore, the fifth parameter δ is significantly different from zero.

5.2 USS Halfbeak diesel engine data


Marciano et al. (2012) used the data to illustrate the application of the McDonald-gamma distribution (Mc- Γ). The distribution of the data is highly skewed to the left and platykurtic (skewness = -1.576 and kurtosis = 1.653). The MLEs (with corresponding standard errors in parentheses) of the parameters of NW{C} distribution and the statistics AIC, the log-likelihood value and K-S and the corresponding p-values are given in Table 3. The values of AIC, log-likelihood and K-S statistics for Mc- Γ , Kumaraswamy-gamma distribution (Kw- Γ) are taken from Marciano et al. (2012). The other results in Table 3 are obtained by using NLMIXED procedure in SAS and the MATLAB software.

The NW{C} distribution has the smallest AIC and K-S statistics, and the largest log-likelihood value, which indicates NW{C} is superior to the other distributions in Table 3. Figure 5 displays the estimated PDF of the NW{C}, Mc- Γ and Kw- Γ

Table 3 Parameter estimate (standard error in parentheses) for USS Halfbeak diesel engine data

	Four-parameter NW{C}	McDonald-gamma	Kumaraswamy-gamma
MLE (SE ^b)	$\hat{\mu} = 6.072 (1.514)$ $\hat{\sigma} = 4.217 (1.207)$ $\hat{c} = 1.448 (0.161)$ $\hat{\gamma} = 10.073 (0.936)$	$\hat{a} = 99.865 (0.294)$ $\hat{\beta} = 2.030 (0.022)$ $\hat{\alpha} = 0.0421 (0.006)$ $\hat{b} = 200.040 (60.178)$ $\hat{c} = 0.2796 (0.014)$	$\hat{a} = 6.384 (0.992)$ $\hat{\beta} = 0.1996 (0.040)$ $\hat{b} = 2.403 (0.001)$ $\hat{c} = 0.0013 (0.000)$
Log-likelihood	-196.45	-217.35	-239.2
AIC	400.9	444.7	486.4
K-S	0.0811	0.2635	0.2849
P-value	0.739	1.045e-04	1.974e-05

b: standard error of the MLE.



distributions. The figure shows that the $NW\{C\}$ distribution provides the best fit to the data compared to other distributions.

6. Conclusions

This paper presents a method to generate the $T-X(W)$ families of distributions introduced in Alzaatreh et al. (2013b) by defining the W function using the quantile function of another random variable Y . Table 1 contains some $T-X\{Y\}$ families based on different quantile functions. The $T-X\{Y\}$ framework provides an easy way for generating distributions of the $T-X(W)$ family introduced by Alzaatreh et al. (2013b). Existing methods like the methods of combination reviewed in Lee et al. (2013) for generating univariate continuous distributions can be derived using the $T-X\{Y\}$ framework. $T-X\{exponential\}$, $T-X\{Weibull\}$ and $T-X\{Rayleigh\}$ can be viewed as families of distributions arising from hazard functions. The $T-X\{Y\}$ family is extended by using survival function of X . The family of life distributions derived by Marshall and Olkin (1997) can be derived using the $T-X\{Y\}$ family based on survival functions. Three new distributions in the family, normal-Weibull{Cauchy}, normal-Weibull{logistic} and Weibull-uniform{log-logistic} distributions are defined. These distributions are very flexible and are capable of fitting various types of data. The Old Faithful Geyser eruption data are used to illustrate that the $NW\{C\}$ distribution fits bimodal data very well, which typically can only be adequately fitted using mixture distributions.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

The authors, viz MAA, CL and FF with the consultation of each other carried out this work and drafted the manuscript together. All authors read and approved the final manuscript.

Acknowledgments

We are grateful for many constructive comments and suggestions from the associate editor and the two referees. These comments and suggestions have greatly improved the presentation of the paper.

Received: 10 December 2013 Accepted: 1 April 2014

Published: 11 June 2014

References

- Akinsete, A, Famoye, F, Lee, C: The beta-Pareto distribution. *Statistics* **42**, 547–563 (2008)
- Alexander, C, Cordeiro, GM, Ortega, EMM, Sarabia, JM: Generalized beta-generated distributions. *Computational statistics and data analysis* **56**(6), 1880–1896 (2012)
- Alshawarbeh, A, Lee, C, Famoye, F: The beta-Cauchy distribution. *Journal of Probability and Statistical Science* **10**, 41–58 (2012)
- Alzaatreh, A, Famoye, F, Lee, C: Gamma-Pareto distribution and its applications. *Journal of Modern Applied Statistical Methods* **11**(1), 78–94 (2012a)
- Alzaatreh, A, Lee, C, Famoye, F: On the discrete analogues of continuous distributions. *Statistical Methodology* **9**, 589–603 (2012b)
- Alzaatreh, A, Famoye, F, Lee, C: Weibull-Pareto distribution and its applications. *Communications in Statistics-Theory and Methods* **42**, 1673–1691 (2013a)
- Alzaatreh, A, Lee, C, Famoye, F: A new method for generating families of continuous distributions. *Metron* **71**(1), 63–79 (2013b)
- Arellano-Valle, RB, Cortés, MA, Gómez, HW: An extension of the epsilon-skew-normal distribution. *Communications in Statistics-Theory and Methods* **39**(5), 912–922 (2010)
- Ascher, H, Feingold, H: *Repairable Systems Reliability*. Marcel Dekker, New York (1984)
- Azzalini, A: A class of distributions which includes the normal ones. *Scand J Stat* **12**, 171–178 (1985)
- Cordeiro, GM, de Castro, M: A new family of generalized distributions. *J Stat Comput Simul* **81**(7), 883–898 (2011)
- Cordeiro, GM, Lemonte, AJ: The β -Birbaum-Saunders distribution: an improved distribution for fatigue life modeling. *Computational Statistics and Data Analysis* **55**(3), 1445–1461 (2011)
- Cordeiro, GM, Ortega, EMM, Nadarajah, S: The Kumaraswamy Weibull distribution with application to failure data. *J Franklin Inst* **347**, 1399–1429 (2010)
- Cordeiro, GM, Nadarajah, S, Ortega, EMM: The Kumaraswamy Gumbel distribution. *Statistical Methods and Applications*, 2012 **21**(2), 139–168 (2011)
- Dekking, FM, Kraaikamp, C, Lopuhaä, HP, Meester, LE: *A Modern Introduction to Probability and Statistics*. Springer, New York (2005)
- Eugene, N, Lee, C, Famoye, F: The beta-normal distribution and its applications. *Communications in Statistics-Theory and Methods* **31**(4), 497–512 (2002)
- Famoye, F, Lee, C, Eugene, N: Beta-normal distribution: bimodality properties and applications. *Journal of Modern Applied Statistical Methods* **3**(1), 85–103 (2004)
- Ferreira, JTAS, Steel, MFJ: A constructive representation of univariate skewed distributions. *J Am Stat Assoc* **101**, 823–829 (2006)
- Härdle, W: *Smoothing Techniques with Implementation in S*. Springer, New York (1991)
- Johnson, NL: Systems of frequency curves generated by methods of translation. *Biometrika* **36**, 149–176 (1949)
- Johnson, NL, Kotz, S, Balakrishnan, N: *Continuous Univariate Distributions*, Vol. 1, 2nd edn. John Wiley and Sons, Inc., New York (1994)
- Jones, MC: Families of distributions arising from distributions of order statistics. *Test* **13**, 1–43 (2004)
- Jones, MC: Kumaraswamy's distribution: a beta-type distribution with tractability advantages. *Statistical Methodology* **6**, 70–81 (2009)
- Kotz, S, Vicari, D: Survey of developments in the theory of continuous skewed distributions. *Metron* **LXIII**, 225–261 (2005)
- Kumaraswamy, P: A generalized probability density functions for double-bounded random processes. *J Hydrol* **46**, 79–88 (1980)
- Lai, CD: Constructions and applications of lifetime distributions. *Appl Stoch Model Bus Ind* **29**, 127–140 (2013)
- Lee, C, Famoye, F, Alzaatreh, A: Methods for generating families of univariate continuous distributions in the recent decades. *WIREs Computational Statistics* **5**, 219–238 (2013)
- Marciano, FWP, Nascimento, ADC, Santos-Neto, M, Cordeiro, GM: The Mc- Γ distribution and its statistical properties: an application to reliability data. *International Journal of Statistics and Probability* **1**(1), 53–71 (2012)
- Marshall, AW, Olkin, I: A new method for adding a parameter to a family of distributions with applications to the exponential and Weibull families. *Biometrika* **84**, 641–652 (1997)
- Marshall, AW, Olkin, I: *Life Distributions*. Springer, New York (2010)
- McDonald, JB: Some generalized functions for the size distribution of income. *Econometrica* **52**, 647–663 (1984)
- McDonald, JB, Xu, YJ: A generalization of the beta distribution with applications. *J Econ* **66**, 133–152 (1995)
- Meeker, WQ, Escobar, LA: *Statistical methods for reliability data*. John Wiley & Sons, New York (1998)
- Pearson, K: Contributions to the mathematical theory of evolution. II. Skew variation in homogeneous material. *Philos Trans R Soc Lond A* **186**, 343–414 (1895)
- Polyanin, AD, Manzhirov, A: *Handbook of Integral Equations*, 2nd edn. Chapman & Hall/CRC, New York (2008)
- Shannon, CE: A mathematical theory of communication. *Bell System Technical Journal* **27**, 379–432 (1948)
- Shorack, GR, Wellner, JA: *Empirical Processes with Applications to Statistics*. John Wiley & Sons, New York (1986)
- Tukey, JW: The Practical Relationship Between the Common Transformations of Percentages of Counts and Amounts. Technical Report 36. Princeton University, Princeton, NJ, Statistical Techniques Research Group (1960)

doi:10.1186/2195-5832-1-2

Cite this article as: Aljarrah et al.: On generating T-X family of distributions using quantile functions. *Journal of Statistical Distributions and Applications* 2014 **1**:2.